10/537421

RECEIVED CENTRAL FAX CENTER NOV 2 0 2008

7. (previously presented) Process for the preparation of aqueous dispersions of non-ionic – N=C=O blocked polyisocyanates according to claim 6., wherein the non-ionic alkoxylated diols (iii) have the general formula I:

$$R_1CH_2O-(-CH_2CH_2O)_n-(CH_2CHCH_3O)_m-R_2$$

(I)

wherein:

and R₂ is methyl, R₃ is ethyl, n is a number from 15 to 30 and m is a number from 0 to 10.

- 8. (previously presented) Process for the preparation of aqueous dispersions of non-ionic –N=C=O blocked polyisocyanates according to claim 6. wherein the polyisocyanate (i) is an isocyanurate obtained from 1,6-hexamethylenediisocyanate and a reaction product of trimethylol propane and toluenediisocyanate.
- 9. (previously presented) Process for the preparation of aqueous dispersions of non-ionic -N=C=O blocked polyisocyanates according to claim 6. wherein the step b. is preceded by dilution of the reaction mixture obtained in a. with from 0.10 to 0.50 parts by weight of a water mixable polar solvent.
- 10. (previously presented) Process for the preparation of aqueous dispersions of non-ionic

- -N=C=O blocked polyisocyanates according to claim 9., wherein the water mixable polar solvent is selected from the group consisting of methyl ethyl ketone, acetone, and cyclohexanone.
- 11. (previously presented) Process for the preparation of aqueous dispersions of non-ionic –N=C=O blocked polyisocyanates according to claim 6., wherein in step a. the equivalent ratio of polyisocyanate (i) and alkoxylated diol (iii) is such that the percentage in weight of the ethoxyl groups is from 20 to 30%.
- 12. (previously presented) Process for the preparation of aqueous dispersions of non-ionic –N=C=O blocked polyisocyanates according to claim 6., wherein the blocking agent (ii) is from the group consisting of butanone oxime and 3,5-dimethylpyrazole.
- 13. (previously presented) Process for the preparation of aqueous dispersions of non-ionic

 -N=C=O blocked polyisocyanates according to claim 6., wherein the amount of blocking agent

 (ii) is such that the equivalent ratio of the isocyanate groups of the oligomer and the blocking agent (ii) is from 1:1 to 1:1.2.
- 14. (previously presented) Process for the preparation of aqueous dispersions of non-ionic –N=C=O blocked polyisocyanates according to claim 6., wherein in step c. the mixture is dispersed into water under stirring to obtain a dispersion having a solid content of from 25 to 35% by weight.
- 15. (previously presented) Procedure for the oil- and/or water-repellent finishing of textiles, characterized by the fact that, as a finishing agent, an aqueous composition is used, said aqueous composition comprising an organic perfluorinated polymeric compounds and from 0.1 to 10% by weight of the total weight of the composition, of an aqueous dispersion of a non-ionic –N=C=O blocked polyisocyanates according to claim 1., the weight ratio between the solid fraction of the

Fax:7132668510

aqueous dispersion and the perfluorinated polymeric organic compounds being from between 1:1 and 1:15.

- A textile printing paste comprising from 0.3 to 5% by weight of an 16. (previously presented) aqueous dispersion of claim 1.
- The textile printing paste of claim 16., wherein the aqueous dispersion 17. (previously presented) of claim 1. is present at a concentration of from 1 to 3.5%.
- Aqueous dispersion according to claim 4., wherein the 18. (previously presented) toluenediisocyanate is composed of 2,4 and 2,6 isomers being in a weight ratio of 80:20.
- Process fro the preparation of aqueous dispersion according to claim 19. (previously presented) 8., wherein the toluenediisocyanate is composed of 2,4 and 2,6 isomers being in a weight ratio of 80:20.